



# Design principles of the bacterial quorum sensing gene networks

Andrew B. Goryachev\*

Bacterial quorum sensing (QS) has attracted much interest as the manifestation of collective behavior in prokaryotic organisms once considered strictly solitary. Significant amount of genetic, biochemical, and structural data which, has been accumulated in studies on QS in many species allows us to map properties of specific molecules and their interactions on the observed population-wide bacterial behavior. The present review attempts to give a systems biology perspective on the structure of genetic regulatory networks that control QS and considers functional implications of a variety of design principles that recur in the organization of these networks across species. © 2009 John Wiley & Sons, Inc. *WIREs Syst Biol Med* 2009 1 45–60

Quorum sensing (QS) refers to the ability of bacterial populations to coordinately regulate gene expression in response to changes in the local population density. Some examples of such collective behavior include secretion of antibiotics,<sup>1–3</sup> production of extracellular matrix,<sup>4–6</sup> emission of light,<sup>7,8</sup> and switch to swimming or swarming motility.<sup>9–11</sup> The cooperativity is achieved through the exchange of signaling molecules that are produced by individual bacterial cells and secreted in the surrounding extracellular space. QS therefore represents an example of a cooperative cell-cell communication<sup>12</sup> that occurs at a distance rather than through a direct physical contact of interacting cells.

Since the discovery of this phenomenon in marine luminescent bacteria *Vibrio fischeri*, a symbiont of a tropical squid, and its free-living relative *Vibrio harveyi*,<sup>13–15</sup> the QS research has evolved into a large and flourishing field that provided a totally new perspective on the general physiology, genetics, ecology, and pathogenicity of microorganisms. Indeed, the discovery of the QS promoted a dramatic shift in the microbiologist's perception of bacteria from dull solitary organisms incapable of any complex cooperative behavior to agile team players who continuously survey their environment for the presence of their kin, competitors, and enemies.

Comprehensive coverage of the vast field of QS, even in a succinct form, is far beyond the

scope of this review. Historic perspective and modern views on the physiology and molecular biology of QS in individual bacterial species as well as extensive comparative analyses across multiple species can be found in many excellent reviews written by the pioneers of this field.<sup>15–39</sup> Burgeoning development in the past decade resulted in branching and diversification of the QS research. Thus, an unanticipated complexity of bacterial communication within the species, between close and distant taxons as well as with eukaryotic organisms (interkingdom signaling),<sup>40</sup> fostered the emergence of a new discipline of bacterial sociobiology that focuses on the evolutionary and ecological aspects of bacterial communication and cooperation.<sup>39,41,42</sup> Another newly formed branch studies QS from the perspective of the host-pathogen interactions<sup>43–46</sup> and seeks to develop novel antibacterial drugs that will interfere with QS and prevent or reduce bacterial pathogenicity toward animal<sup>47–49</sup> and plant<sup>45,50</sup> hosts.

The present contribution attempts to give a systems biology perspective on the intracellular mechanisms that regulate QS behavior on the molecular level. Unicellular nature, genetic tractability, and availability of complete genomic sequences have long placed bacterial regulatory networks into the focus of systems biology.<sup>51–54</sup> The QS phenomenon in particular provides an unprecedented opportunity to study molecular origins of collective multicellular behavior and perform comparative analyses of distinct network designs which evolved under the pressure of diverse ecological requirements.

\*Correspondence to: Andrew.Goryachev@ed.ac.uk

Centre for Systems Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, UK

DOI: 10.1002/wsbm.027

## WHAT IS 'QUORUM SENSING NETWORK'?

I refer to quorum sensing network (QSN) as a network of signaling and gene regulation events that are responsible for the production and perception of the signaling molecules as well as for the final outcome of communication—regulation of target genes whose products directly control specific functions, e.g., encode bacterial flagella. Depending on the complexity of the network, several layers of regulatory molecules may intercalate between the signaling molecules, also known as *autoinducers* (AIs), and the QS transcription factors (QSTRs) responsible for regulation of the target genes. Practically in all studied cases the QSN does not stand on its own but is instead imbedded into a complex regulatory network which integrates multiple environmental stimuli, such as pH, temperature, and the availability of nutrients.

Typically, the QSN consists of multiple components which can be placed into several functional categories. (1) Enzymes that produce signaling molecules, AI synthases, and the respective AI secretion systems. (2) QS sensor systems and transduction pathways that connect the receptors for signaling molecules to the regulators of QSTRs and possibly integrate other sensory inputs. (3) QSTRs and their direct regulators—proteins, RNA species, or metabolites that directly affect the activity of QSTRs themselves or regulate their transcription and translation.

## QS SIGNALING MOLECULES, THEIR SYNTHASES AND SENSOR SYSTEMS

The choice of a signaling molecule, the physical messenger of cell–cell communication, imposes specific requirements on the presence and molecular properties of the components of the QSN that directly interact with the messenger. Indeed, membrane permeability, diffusivity, and stability to nonenzymatic degradation in the environment are all important biophysical properties of a signaling molecule that specify its signaling range and the means of its secretion and detection. Historically, two QS-signaling systems came to the limelight first: *N*-acylhomoserine lactones (AHLs) of Gram-negative bacteria<sup>16,23,55,56</sup> and peptides of Gram-positive species.<sup>57–60</sup> Early studies suggested that AHLs are low-molecular-weight metabolites that can passively diffuse through the cellular membranes. Many of these AIs were shown to bind directly to the QSTRs influencing their oligomerization state and affinity to the DNA *cis*-regulatory sequences.<sup>55</sup> Within this early conceptual model, AHLs require neither special secretory machinery nor cell-surface receptors. In

contrast, peptides of Gram(+) bacteria are actively extruded from the cells and detected by transmembrane receptors<sup>61</sup> or actively imported back into the cells.<sup>62,63</sup> Further studies, however, reduced the divide between the two QS systems and identified a number of common features. Thus, it was shown that some AHLs with long acyl chains can hardly penetrate the membrane and thus require assisted transport.<sup>64</sup> Moreover, in some Gram(–) bacteria, e.g., *Vibrio* species,<sup>35</sup> AHLs are detected by cell-surface receptor kinases in a Gram(+) fashion.

AHLs are synthesized from metabolites readily produced in bacterial cells.<sup>26,27</sup> The best characterized family of AHL-synthases are homologs of *V. fischeri* LuxI, which are usually encoded in pairs with matching LuxR-type QSTRs.<sup>16,65,66</sup> Other, non-LuxI-type synthases, such as AinS (*V. fischeri*) and LuxM (*V. harveyi*),<sup>67</sup> with biochemical properties both similar and distinct from LuxI-type enzymes<sup>56</sup> have also been described. The length of the acyl chain is an important property of an AHL molecule. Indeed, chains shorter than four result in highly chemically unstable AHLs, which renders them generally unsuitable for cell–cell communication,<sup>38</sup> while AHLs with chains longer than 10 poorly diffuse through the cell membranes and might require transporters for their excretion.<sup>64</sup> Often, a single AHL synthase can produce a broad repertoire of AHLs with varying lengths of acyl chains.<sup>68</sup> Importantly, bacterial genomes also encode enzymes that degrade AHLs. Regulated activation of these enzymes, for example, in response to a stress signal ppGpp, can efficiently abrogate QS signaling.<sup>69,70</sup>

Peptide messages of Gram(+) species are genetically encoded as precursor peptides and do not require synthases *per se*.<sup>60,61,71</sup> Instead, these bacteria encode transmembrane enzymes, which proteolytically process the precursors while extruding them into the extracellular space. These proteins thus serve the function of synthases and exporters simultaneously. Two strategies are in use for detection of peptides in the environment: receptor histidine kinases<sup>61</sup> and active importers, usually from the family of ABC transporters.<sup>62</sup> Once imported into the cell, the peptides are sensed by intracellular receptors, such as *Bacillus subtilis* phosphatase Rap, whose activity is repressed by peptide binding,<sup>72</sup> thus preventing it from dephosphorylating a transcriptionally active response regulator.<sup>62</sup>

Both Gram-positive and Gram-negative species utilize a variety of furanones, collectively known as AI-2.<sup>30</sup> AI-2 molecules are a result of spontaneous cyclization of 4,5-dihydroxy-2,3-pentanedione (DPD), which together with homocysteine are the products of the LuxS group of enzymes first described

in *V. harveyi* and presently found in over 50 bacterial species.<sup>21,30,73,74</sup> Sensing of AI-2 as a QS signal has been studied in detail in *V. harveyi* and other marine vibrios in which AI-2 is bound by periplasmic protein LuxP and this complex is sensed by the transmembrane receptor LuxQ.<sup>75</sup> A different mode of interaction, which involves AI-2 internalization and degradation, has been described in enteric bacteria.<sup>76</sup> In these species, AI-2 is phosphorylated by intracellular kinase LsrK. AI-2-P then binds to and deactivates transcriptional repressor LsrR, which represses the *lsr* operon. First four genes of this operon (*lsrA*–*lsrD*) encode an AI-2-specific ABC importer, while products of the following genes, e.g., *lsrG*, are involved in the subsequent degradation of AI-2-P. Appearance of AI-2 in the extracellular medium thus first causes its low-level and possibly nonspecific internalization that activates the *lsr* operon, this in turn causes specific import and degradation of the messenger.

Expansion of the QS research resulted in the discovery of other signaling molecules with diverse chemical nature. In addition to AHLs, Gram(–) bacteria were also found to produce alkyl-quinolones,<sup>77,78</sup> DSF (*cis*-11-methyl-2-dodecenoic acid),<sup>79</sup> and fatty acid methyl esters.<sup>80</sup> Recently, the nature of an elusive *Vibrio cholera* AI CAI-1 has been identified and found to belong to yet another class of chemical compounds, 3-hydroxytridecan-4-one.<sup>81</sup> Not restricted to peptide signaling only, some Gram(+) bacteria, e.g., streptomycetes, also synthesize  $\gamma$ -butyrolactones.<sup>82</sup> Future screens designed to identify bacterial cell–cell communication mediators will likely reveal additional signaling molecules.

## NETWORK CORE: THE LUXR/LUXI PARADIGM IN GRAM-NEGATIVE BACTERIA

Already upon their original identification,<sup>14</sup> QSTR LuxR and the respective AI synthase LuxI were shown to be essential proteins for the QS regulation of luminescence in *V. fischeri*. A regulatory subnetwork based on the two genes, their protein products, and LuxI-generated AI rapidly became a paradigm for the QSN organization in Gram(–) bacteria.<sup>16</sup> Phylogenetic analyses of many genes encoding LuxR/LuxI homologs in various species<sup>65</sup> indicated that these genes coevolved as mutually regulating couples. This justifies their contemporary consideration as core regulatory modules of the AHL-based QSNs. Moreover, phylogenetic analysis shows that *luxR/luxI* homologs fall into two distinct classes.<sup>83</sup> Class A systems are widespread among various divisions of proteobacteria and their

LuxR-type proteins are typically transcription activators. Class B systems, identified so far only in  $\gamma$ -proteobacteria, show a distinct pattern of genomic organization and their QSTRs are mostly repressors. LuxR-type QSTRs of both classes possess N-terminal AI-binding and C-terminal DNA-binding domains. Binding of cognate AIs to the LuxR-type proteins in stoichiometry 1 : 1 may result in either increase (class A) or attenuation (class B) of binding to the respective DNA *cis*-elements. Most of the class A QSTRs, such as the prototypical *V. fischeri* LuxR itself, are activators that require their cognate AIs for binding to DNA and transcriptional activation.<sup>27,83</sup> Competence for DNA binding in most of these QSTRs is correlated with their oligomerization state. Typically, these QSTRs are monomers in the inactive AHL-free state but dimerize efficiently in response to the AHL binding. This dimerization might be responsible for the observed protein half-life extension of the AHL-bound QSTRs.<sup>84–86</sup> Once dimeric, they strongly and specifically bind to the inverse-repeat DNA sequences of about 20 bp known as *lux-type boxes*.<sup>16</sup> Some exclusions from the above rules have also been reported. Thus *Erwinia caratovora* CarR was shown to be dimeric and competent to bind DNA already in the AHL-free state and formed higher-order oligomers in response to the AHL addition.<sup>87</sup>

More recently, a number of LuxR-type repressors, whose transcription repression is relieved by binding to AHLs, has been described.<sup>88–92</sup> Proteins of this type have been shown to be dimeric already in the AHL-free state,<sup>90</sup> which is consistent with their ability to bind palindromic DNA sequences and repress target genes in the absence of AHLs. Binding of AHLs to QSTRs triggers a conformational switch to the state that can no longer bind DNA. The complex releases DNA and transcription of target genes is activated. Interestingly, in both activator and repressor cases, accumulation of AIs effectively results in activation of the target genes.

While the ability of LuxR-type QSTRs to increase or reduce their affinity for DNA in response to AI binding appears to be a fundamental molecular property, their activator or repressor function should depend on the position of the respective DNA *cis*-element within the upstream regulatory sequence of a specific gene. Thus, conceivably the same QSTR could be activator for some and repressor for other target genes. Despite this theoretical possibility, class A QSTRs have been largely characterized as activators, while those of class B as repressors. ExpR1 and ExpR2 QSTRs recently characterized in *E. caratovora* may present an example of an alternative behavior.<sup>93,94</sup> In the AHL-free state, both proteins are DNA-bound and

possess other properties of class B QSTRs. However, they both activate transcription of regulatory mRNA-binding protein RsmA. 3-Oxo-C8-HSL selectively binds to ExpR2 and releases it from DNA, thus partially attenuating *rsmA* transcription. Another *E. caratovora*-produced AHL, 3-oxo-C6-HSL, binds to both QSTRs and releases them from DNA, further reducing *rsmA* transcription. Ironically, since RsmA is a repressor of the target genes (plant cell wall degrading enzymes), transition to the high-density state ('transition to quorum') results in the activation of target genes also in this case.

Importantly, *luxR/luxI* modules of different classes exhibit distinct genomic organization and regulatory interactions. Class A activators are presently understood the best. As shown schematically in Figure 1(a), a LuxR homolog is typically encoded separately from the operons it regulates and *luxI* is transcribed divergently or in the same direction as the *luxR* homolog. Normally, expression of the *luxI* homolog is positively regulated by the LuxR-type QSTRs.<sup>16</sup> Provided that LuxI-type synthases generate AI molecules necessary for the activity of the QSTRs, this activation of *luxI* transcription by LuxR constitutes the classical QS positive feedback loop that amplifies the intracellular production of AI in response to the extracellular AI. Although in this case production of the monomeric transcription factor remains at the same level, the copy number of the mature dimeric QSTR increases due to the increased formation of the LuxR-AI complex. Additional positive feedback loop may be provided by autoactivation of *luxR* transcription. This autoregulation can also amplify the copy number of the monomeric transcription factor and has been demonstrated at least for some QSTRs, e.g., for the prototypical *V. fischeri luxR*<sup>95</sup> and *E. caratovora carR*.<sup>87</sup>


In contrast, class B *luxR/luxI* modules of  $\gamma$ -proteobacteria exhibit strikingly different patterns of genomic organization and coregulation (Figure 1(b)). Similar to the described first *esaR/esaI* module,<sup>88,90</sup>

other characterized representatives of this class (e.g., *expR/expI* of *E. chrysanthemi*<sup>92</sup> and *ypsR/ypsI* of *Y. pseudotuberculosis*<sup>96,97</sup>) are typically transcribed convergently with a small overlap between the transcribed frames of 20–25 bp, significance of which is not presently understood.<sup>89,96</sup> Unlike in class A, transcription of the *luxI* homolog generally is not directly regulated by the LuxR homolog and is controlled by other regulators.<sup>97</sup> As the QSTRs of this class are normally transcriptional repressors, autoinhibition of the *luxR* transcription is not uncommon but not obligate.<sup>90,97</sup>

In all known systems, mutual regulation within the *luxR/luxI* module accounts for only a part of the overall transcription regulation of the *luxR* and *luxI* homologs. Both may possess constitutive promoters as well as promoters regulated by nutrients,<sup>98</sup> signaling molecules,<sup>99</sup> and upstream transcription factors including other QSTRs.<sup>97,100</sup>

## OTHER TYPES OF QSTRS

While characterized best, *luxR/luxI* paradigm does not exhaust all possibilities in bacterial QSNs. Even in Gram(–) species, *V. fischeri* LuxR homologs are not the only known QSTRs. Ironically, the best characterized examples of alternative QSTRs have been described in Vibrios, the same genus that gave birth to the *luxR/luxI* paradigm. Starting with the luminescence regulator LuxR in *V. harveyi*<sup>101</sup> (not related to *V. fischeri* LuxR, the protagonist of the LuxR class of QSTRs), a number of dimeric transcription factors that do not require AI molecules to bind to DNA were identified. Other representatives of this class of QSTRs are HapR in *V. cholera*<sup>102–104</sup> and LitR in *V. fischeri*.<sup>105</sup> Sequence and structure<sup>106</sup> homology indicate that these QSTRs belong to a large class of TetR transcription factors, dimeric proteins that bind palindromic *cis*-regulatory DNA sequences.<sup>107</sup> The protagonist of the class,



**FIGURE 1** | Genomic organization and mutual regulation within the QSN core module. (a) Class A *luxR/luxI* module of Gram(–) bacteria. (b) Class B *luxR/luxI* module. (c) Representative example of Gram(+) QSN core module. Transcription regulator AgrA activates transcription of the *agr* operon. *agrD* and *agrB* encode AI peptide precursor and peptide extruding enzyme respectively. Two strands of DNA are represented by separate lines. Open boxes symbolize *cis*-regulatory elements. Regulatory interactions are shown by hammerheads (negative) and arrows (positive). Interactions found only in a subset of characterized systems are drawn by dashed line.

transcription repressor TetR controls the tetracycline exporter *tetA*, whose expression is activated when tetracycline binds to the TetR dimer and TetR dissociates from the *tetA* promoter.<sup>107</sup> Intriguingly, HapR, whose structure was recently determined,<sup>106</sup> also has a putative ligand-binding domain. Although TetR factors are generally considered as transcription repressors, their actual mode of action depends on the specific gene. Thus, HapR, most known for its repression of the *V. cholera* virulence master-regulator *aphA*, is also an activator of hemagglutinin HA/protease<sup>102</sup> and DNA uptake regulator.<sup>108</sup> As many other TetR transcription factors, HapR also represses its own expression.<sup>109</sup> The QS regulation of this class of QSTRs is believed to be performed by sRNA species, which effectively annihilate QSTR mRNA at low population density<sup>110–112</sup> (see a detailed discussion below).

QSTRs of Gram(+) bacteria, which are usually represented by the response regulators of two-component sensor systems, are typically found to positively regulate themselves and the ORFs encoding peptide precursors and their exporters.<sup>61</sup> This ensures that copy numbers of both the QSTRs and the signaling molecules are amplified in response to the QS signal. Genomic organization of Gram(+) QSN core modules is species specific.<sup>58</sup> The QSTR, signaling peptides, their receptors, and exporters may be encoded by the same operon, as in the case of *agr* operon of *Staphylococcus aureus*<sup>60,113</sup> (Figure 1(c)), or split into several operons co-regulated by the phosphorylated QSTR.<sup>58,61</sup>


## IS THERE A SWITCH IN QSN AND WHAT FLIPS IT?

Principles that underlie operation of the QSN have captivated attention of researchers since the discovery of the QS phenomenon. The main function of the QSN, to control the copy number of the QSTR in response to the changing extracellular concentration of AI, can be achieved in two distinct ways. The QSN may operate as a rheostat by gradually changing the copy number of the QSTR and thus the expression of target genes. Alternatively, it may operate as a bistable switch with two distinct 'on' and 'off' states.<sup>114,115</sup> In the latter situation, an individual cell would be likely found in either uninduced or fully induced states, while cells with intermediate expression levels of the QSTR would be extremely rare.<sup>116,117</sup>

A well-known fact in mathematical sciences that positive feedback is a necessary condition for multistability<sup>115,118</sup> prompted a hypothesis that the core *luxR/luxI* module may, in fact, implement

such a bistable switch. Assuming that the mature QSTR is a dimer that activates its own transcription, mathematical modeling of a *luxR/luxI*-type module demonstrated that, in principle, this network layout alone is sufficient to provide a bistable switching behavior.<sup>119,120</sup> Further analysis performed with realistic values of network reaction rates<sup>121</sup> identified a number of constraints imposed on this QSN layout and suggested its poor performance as a gene expression switch under the conditions of molecular noise.<sup>122,123</sup> Importantly, the *luxR/luxI* module may function as an autonomous, self-sufficient switch only if the QSTR is a relatively unstable dimer and the binding between the AI and the LuxR-type monomer is not very strong.<sup>121</sup> While this assumption implicitly made in Refs [119,120] appears to hold for some QS systems,<sup>84,124</sup> it is not universal for all *luxR/luxI* modules. Thus, *Agrobacterium tumefaciens* QSTR TraR forms practically irreversible complex with its cognate AI.<sup>125</sup> In fact, the molecule of AHL was found so deeply buried into the protein 3D structure that it was suggested that the complex forms only during TraR translation. Translated in the absence of AHL, TraR is misfolded, poorly soluble, and is rapidly degraded.<sup>85,86,126</sup>

Recent analyses of QS in *V. harveyi* and *V. cholera* suggested an alternative mechanism that can provide robust switchlike behavior without bistability.<sup>110</sup> In both species, AI signals are transmitted to the transcription regulator LuxO, which is phosphorylated in the absence of AIs (low cell density). Phosphorylated LuxO activates transcription of several Qrr small regulatory RNAs. In cooperation with chaperone Hfq,<sup>127</sup> these sRNAs form stable complexes with mRNAs of QSTRs LuxR (*V. harveyi*) or HapR (*V. cholera*). The Qrr sRNAs sequester QSTR mRNA and, either through the formation of an untranslatable complex or through the accelerated mRNA degradation, effectively prevent its translation. As opposed to repression by transcription factors, this type of post-transcriptional regulation results in the destruction of both the target (QSTR mRNA) and the regulator (Qrr sRNA) (Figure 2(a)). The target and regulatory RNAs mutually titrate each other and the outcome of their competition is completely defined by their respective copy numbers, which in turn sensitively depend on the rates of transcription. If transcription rate of either regulator or target (potentially both) depends on the concentration of AI, this mutually antagonistic interaction between RNA species may result in an ultrasensitive response<sup>128,129</sup> of the QSTR copy number to a gradual change in the AI concentration.<sup>110</sup> Thus, rapid, switchlike transition to quorum can be, in principle, achieved in a narrow



**FIGURE 2** | Regulation of QSNs by mutual inhibition. (a) RNA–RNA switch in the QSN of *V. harveyi*. (b) RNA–RNA switch in the QSN of *S. aureus*. The QS positive feedback loop of *agr* operon is shown schematically. (c) Protein–protein mutual inhibition operates as a switch in *A. tumefaciens* QSN. AAI is Agrobacterium AI. S symbolizes substrates of AI synthase TraI. (d) Typical ultrasensitive behavior for expression of the QSTR and its inhibitory regulator schematically represented as a function of the extracellular concentration of AI. Small open boxes represent complex formation.  $\emptyset$  symbolizes eventual degradation of the inhibitory complex. RNA species are shown as rectangles and proteins as ovals. Translation of mRNA species is shown by open arrowheads.

range of changing concentration of the extracellular AI (schematically illustrated in Figure 2(d)).

The RNA-based mutual-antagonism switching mechanism has been also reported in other QSNs. Perhaps the oldest known example of this type of regulation is provided by RNAIII regulatory sRNA of *S. aureus agr* system.<sup>130</sup> RNAIII sequesters and inhibits translation of mRNAs for transcription factor Rot and protein A using almost the same mechanism as *Vibrio Qrr* sRNAs (Figure 2(b)). Presumably, activation of *agr* QSN network by its cognate AI peptide results in a similar ultrasensitive regulation of Rot. In addition, RNAIII forms an activatory complex with *hla* mRNA, which encodes  $\alpha$ -hemolysin. Given that *hla* is repressed by Rot, the transition to quorum may result in a sharp switch between *rot* and *hla*.<sup>58</sup> Post-transcriptional control by small RNAs has recently become a major theme in prokaryotic gene regulation<sup>131,132</sup> and has attracted vivid interest among experimentalists and theoreticians alike.<sup>133–136</sup> Of particular prominence in the context of QSNs<sup>137</sup> is the regulatory motif based on the post-transcriptional regulator known as CsrA (in *E. coli*) or RsmA (in *Pseudomonas aeruginosa*, *E. carotovora*, and others) and the small RNA CsrB (RsmB).<sup>132</sup> The homodimeric CsrA/RsmA-type proteins mostly act as inhibitors of target genes by sequestering their mRNAs. Transcription of CsrB/RsmB sRNAs, which can bind simultaneously up to nine CsrA dimers,<sup>132</sup> results in fast and efficient sequestration of CsrA/RsmA repressors and activation of the target genes. In *E. carotovora*, repressor of plant cell wall degrading

enzymes RsmA is under direct transcriptional control of QSTRs ExpR1 and ExpR2, and the respective sRNA RsmB is regulated by a GacS–GacA two-component phosphorelay activated in response to an as yet unknown extracellular signal.<sup>138</sup>

Mutual-antagonism switch of the QSN, however, does not have to depend on sRNAs. It has been long known that the functionality of *A. tumefaciens* QSN critically depends on the antiactivator TraM.<sup>139,140</sup> Knocking this protein out resulted in a constitutively high copy number of the QSTR TraR and a lockup of the network in the high-density, ‘quorum’, state. Recent work has demonstrated that TraM, whose expression is activated by the TraR dimer, is a homodimeric protein that forms highly stable protein complexes with TraR (two TraR dimers : two TraM dimers) and effectively inactivates the QSTR through the formation of this transcriptionally inactive complex<sup>141</sup> (Figure 2(c)). Mathematical analysis of the *A. tumefaciens* QSN<sup>142</sup> confirmed that TraR and TraM form a mutually antagonistic protein couple and demonstrated that they can produce an ultrasensitive switchlike transition to quorum in response to the AI signal (Figure 2(d)).

## INTEGRATION OF QUORUM SENSING INTO GLOBAL GENE NETWORKS

Recent advance in understanding of bacterial gene networks demonstrated that the QSNs receive multiple regulatory inputs from various modules as well as co-regulate many groups of target genes

together with other networks and motifs.<sup>143</sup> The complexity of the regulatory cross-talk between the QSNs and other elements of the cellular control network varies from relatively low, such as in the well-understood case of *traR/traI* QSN that controls conjugal transfer of *A. tumefaciens* Ti plasmids,<sup>98</sup> to exceptionally high, such as in the QS circuitry of *P. aeruginosa*.<sup>100,144,145</sup>


If bacterium possesses several QSNs, they may cooperate following a variety of design patterns. Thus, several well-characterized examples of parallel QSN connection have been described in *Vibrios*.<sup>35</sup> In *V. harveyi*, at least three QS systems operate in parallel<sup>111</sup>: an AHL molecule HAI-1 and two interspecies signals CAI-1 and AI-2. The three signaling molecules are detected by their respective dedicated receptor histidine kinases LuxN, CqsS, and LuxQ, respectively. However, downstream of these receptors, the three signals come together as the common phosphotransferase protein LuxU collects the phosphates produced by the individual receptors and transfers them to the common response regulator LuxO, which in turn controls transcription of the small regulatory RNAs. Similar organization has been reported for *V. anguillarum* QSN,<sup>146</sup> while *V. cholera* seems to have two parallel QSN inputs mediated by generic signals CAI-1 and AI-2.<sup>147</sup> It has been reported that different AI inputs of the parallel QSN of *V. harveyi* translate into distinct levels of the master QSTR LuxR<sup>148</sup>; however, the mechanisms underlying this specificity are not yet completely understood.

A classical example of sequential arrangement of QSNs is provided by *P. aeruginosa* in which the *lasR/lasI* system has been shown to activate the *rhlR/rhlI* QS core module.<sup>149,150</sup> Another instructive example of hierarchical QSN organization has been described in *V. fischeri*, veteran of the QS field.<sup>35</sup> Recent research has demonstrated that, similar to other *Vibrios*, the sRNA-based *ain* QS system is the primary QSN also in this organism.<sup>151</sup> Receptor kinases detect at least two parallel inputs of the *ain* QSN—AinS-produced C8-HSL and AI-2—and through a LuxU-LuxO type phosphorelay control expression of LitR, a homologue of *V. harveyi* LuxR. LitR is thought to regulate the early phases of squid colonization, e.g., repress bacterial motility, and positively regulate the *luxR/luxI* QSN.<sup>151,152</sup> Full activation of the *luxR/luxI* QSN, which requires higher population density for the transition to quorum ( $> 10^{10}$  cells/ml), results in the maximal induction of the *lux* operon as well as activation of at least 18 other genes.<sup>153</sup>

As microbiologists characterize more regulatory network motifs, it becomes increasingly clear that

bacteria utilize essentially all of them in various combinations with the QSNs to achieve maximal flexibility in the control of gene expression. Rcs phosphorelay system,<sup>154</sup> described originally in *Escherichia coli*, provides an example of such a motif. Receptor kinase RcsC relays phosphate to the response regulator RcsB by means of the phosphotransfer protein RcsD. While some genes can be directly regulated by the RcsB homodimer, others require a cofactor RcsA, which forms a heterodimer with RcsB.<sup>154</sup> In *E. amylovora* and *Pantoea stewartii*, the *cps* operon responsible for the synthesis of capsule exopolysaccharide is under the control of the RcsA–RcsB heterodimer.<sup>155</sup> Recently, it has been shown that in *P. stewartii* QS controls capsule production through the regulation of RcsA.<sup>4,156</sup> In an ingenious design, the prototypical repressor EsaR binds to a site between the weak constitutive promoter of RcsA and the strong autoinducible promoter, as shown in Figure 3(a). At high population density, the EsaR-mediated repression is relieved and massive quantities of RcsA are produced in a positive feedback loop.<sup>156</sup> As RcsA–RcsB also represses the flagella production, transition to quorum results in a switch between motile and sessile, capsule protected, life styles. Presumably, since RcsB is directly regulated by the uncharacterized ligand(s) of the RcsC sensor kinase, this integration design allows for the detection of coincidence (or logical AND operator) between QS and specific extracellular stimuli. This control mode, however, remains to be tested experimentally. Boolean AND operator between QS and another extracellular stimulus can be also achieved by hierarchical expression control as exemplified by *A. tumefaciens* QSN. The requirement for the presence of a specific plant-produced nutrient, octopine, is fulfilled by placing the QSTR *traR* within the opine-activated operon *occ*.<sup>125</sup>

Another motif frequently found together with QSNs is the RsmA/RsmB protein–RNA regulatory system, which has been already discussed in this review. In *E. caratovora*, sRNA RsmB is under the control of a GacS–GacA two-component system, while RsmA is positively regulated by ExpR1 and ExpR2.<sup>93,94</sup> When quorum is reached, both ExpR1 and ExpR2 are removed from the DNA and the expression of virulence factors, which were repressed in the absence of AHLs by RsmA, is activated. The same result could have been achieved, in principle, by an unidentified ligand of GacS through the activation of RsmB transcription. Thus, it can be hypothesized that the resulting QSN integration layout implements a logical OR operator allowing activation of the target genes either in response to the unknown signal or upon reaching quorum (Figure 3(b)).



**FIGURE 3** | Integration of quorum sensing with other sensory inputs. (a) Boolean AND logical circuit in the regulatory network of *P. stewartii*. (b) Boolean OR in *E. carotovora*. RNA species, except RsmB, are omitted for brevity. Notations are the same as in Figure 2.

A recent addition to the theme of signal integration is the recognition that the QSNs participate in the regulation of various cellular trends through the pervasive bacterial second messenger *c*-di-GMP.<sup>157–159</sup> Thus, *c*-di-GMP has been shown to positively regulate the formation of biofilm in a number of species including *V. cholera*.<sup>160</sup> In a recent study,<sup>161</sup> Bassler and colleagues demonstrated that in *V. cholera* the QSTR HapR regulates biofilm formation through two parallel pathways. Firstly, HapR binds to and directly represses the expression of the biofilm master regulator *vpsT*. In addition, HapR regulates a number of genes encoding proteins that generate and destroy *c*-di-GMP. The net result of this regulation is the decreased production of *c*-di-GMP in the state of quorum and, thus, downregulation of biofilm formation. While this exciting new story only begins to develop, it is already clear that QSNs, together with other extracellular-regulated networks, are involved in the *c*-di-GMP-dependent gene regulation in many bacterial species.<sup>34,79,162–164</sup>


## CONCLUSION

QS gives bacteria a glimpse of the complex surrounding environment including the information on the presence of other cells of the same and distinct species. The QS detection mechanism is based on the production and reception of diffusible chemical molecules. Local concentration of the AI(s) in the immediate vicinity of the cell is the only actual extracellular input for the decision-making QSN. Given this information, a bacterium cannot distinguish between the situations when it is surrounded by a large number of other AI-producing cells or when it is enclosed alone in a small

diffusion-impermeable compartment.<sup>165,166</sup> Owing to this inevitable ambiguity, the precise function of QS should be interpreted within the context of a specific ecological niche typical to the life style of the species in study.

Generalizing the design of a few currently well-characterized QSNs, it is tempting to hypothesize that all QSNs probably contain some form of a switch based on RNA–RNA, RNA–protein, or protein–protein mutual-antagonism interaction, which may or may not ensure the bistability of the whole network. The classical QS positive feedback loop implemented in all QSNs in one or another way amplifies the extracellular AI signal and likely trips the switch rather than encompasses the switch itself. Despite some theoretical arguments in favor of bistability, whether the QSNs are mostly ‘rheostats’ or ‘switches’ still remains to be determined experimentally. A recent test of this hypothesis in *V. harveyi*<sup>148</sup> resulted in a conclusion suggesting that its QSN operates in a ‘rheostat’ mode. However, the possibility that some of the QSNs exhibit genuinely bistable behavior still remains open.

QSNs are typically integrated with other bacterial sensory/decision-making networks to achieve precise and flexible control of gene expression. Several network motifs found ubiquitously across species enable bacteria to combine the QS signal with other signaling inputs using all three basic logic operators (AND, OR, and NOT), which are necessary to build Boolean decision rules of any desired degree of complexity. This opens a potential for cellular differentiation based on the spatial variation of environmental conditions throughout the population habitat, e.g. within a biofilm.<sup>167–173</sup> Indeed, the relationship between QS and bacterial multicellular



**FIGURE 4** | Behavior of a hypothetical *luxR/luxI* type QSN. (a) Network layout. Notations are the same as in Figure 2. (b) Stationary states of the network at three values of extracellular AI: a—0 nM; b—12 nM; c—24 nM. (c) After removal of the QSTR dimerization from the network layout, only one stationary state remains. (Same values of extracellular AI.) Filled circles and squares represent stable states of the network. Open circle denotes an unstable state of the bistable QSN. (Reaction rate values are adopted from Ref. 121).

morphogenesis has recently become an area of rapid expansion.<sup>39,160,161,174–177</sup> Future research will result in a major progress in understanding of the molecular mechanisms of QS and the connections between the QSN designs and the functional role of QS in the bacterial life style.

### BOX1: MATHEMATICAL MODELING OF THE QSN DEMYSTIFIED

Mathematical modeling has become a powerful tool of biological research and its area of application continues to expand.<sup>118,182</sup> Even a relatively simple model that describes temporary dynamics of the QSN in terms of ordinary differential equations can shed light on the principles of its operation and reveal modes of behavior that are *possible* for a given network design. Conversely, it may also show that some experimentally observed behavior cannot be achieved by a hypothesized network layout and thus additional experimental work is warranted to identify missing or incorrectly characterized network elements.

To describe mathematically the dynamics of a hypothetical QSN shown in Figure 4(a) (generic layout of the *luxR/luxI* type module, adopted from Ref. [121]), one needs to describe how concentrations of the involved proteins, RNA species, and AI change in time. This can be done by using the mass-action rate law of chemical kinetics adapted to model complex cellular processes, such as transcription and translation.<sup>119,120,183,184</sup> For example, we can describe the rate of change for the concentration of *luxR*-like mRNA *r* as follows:

$$\frac{dr}{dt} = k_1 - k_2r.$$

Here, according to the network layout, it is assumed that mRNA *r* is transcribed and degraded constitutively with rates  $k_1$  and  $k_2$  respectively. The mRNA *r* is then translated into a monomeric protein *R*, which reversibly binds a molecule of AI *A* (index *i* indicates intracellular concentration) to form complex *P*, which in turn reversibly dimerizes into a mature QSTR *D*. Describing QSTR-activated transcription of *I* (LuxI-like AI synthase), we assume that the RNA polymerase is abundant, while the availability of activator *D* is the rate limiting factor. However, *I* promoter has a maximal rate of transcription initiation and excess of *D* cannot speed up the transcription past this maximal rate. These requirements can be captured by the Michaelis–Menten kinetic law and the resulting rate of change for the mRNA concentration *i* becomes

$$\frac{di}{dt} = \frac{k_3D}{K_4 + D} - k_5i.$$

Assuming abundance of ribosomes, translation of mRNA can be represented as a simple first-order reaction with a rate proportional to the mRNA concentration. Taking *R* protein concentration *R* as an example, we can describe the rate of *R* change as follows:

$$\frac{dR}{dt} = k_6r - k_7R - k_8R \cdot A_i + k_{-8}P.$$

The four terms in the above equation represent mRNA translation, protein degradation, formation of complex *P*, and its dissociation, respectively. Likewise, for the concentration of AI, which is continuously produced by LuxI, the rate of concentration change

is as follows:

$$\frac{dA_i}{dt} = k_{11}I - k_8R \cdot A_i + k_{-8}P + k_{14}(A_e - A_i),$$

where the last term represents passive diffusive exchange between intra- and extracellular molecules of AI. The latter serves as a free parameter of the model whose variation dictates the entire intracellular dynamics of the QSN.

Once the complete model is constructed (in this case, it consists of seven equations), it can be analyzed numerically on the computer. However, in some simple cases, like the example considered, the behavior of the model can be also studied analytically, e.g. by reducing the number of variables.<sup>119,121,142</sup> Thus, applying standard methods of chemical kinetics it can be shown<sup>121</sup> that  $dA_i/dt$  is a function of  $A_i$ , as shown in Figure 4(b) at three different values of  $A_e$ , the extracellular concentration of AI. Intersections of the

function with zero axis symbolize stationary states of the QSN. The S-shaped form of the function ensures that at some intermediate concentration values of the extracellular AI the network is bistable. Indeed, it simultaneously has two steady states with low, corresponding to the ‘off’ state, and high, representing the ‘on’ state, concentrations of the intracellular AI. In biological terms, this mathematical result can be interpreted so that the given QSN layout *may* possess bistable switchlike behavior at some biologically plausible reaction parameters. If, however, we remove the dimerization of the QSTR from the reaction scheme in Figure 4(a) and instead assume that P is the transcriptionally competent QSTR, we will obtain a very different dependence of  $dA_i/dt$  on  $A_i$  (see Figure 4(c)). This modified QSN layout has a single steady state at any  $A_e$  and with any combination of model parameters. This implies that the variant of the QSN layout without QSTR dimerization *cannot* be a ‘switch’ in a strict sense but instead will always behave as a ‘rheostat’.

## REFERENCES

- Barnard AM, Bowden SD, Burr T, Coulthurst SJ, Monson RE, et al. Quorum sensing, virulence and secondary metabolite production in plant soft-rotting bacteria. *Philos Trans R Soc Lond B Biol Sci* 2007, 362:1165–1183.
- Wei JR, Lai HC. N-acylhomoserine lactone-dependent cell-to-cell communication and social behavior in the genus *Serratia*. *Int J Med Microbiol* 2006, 296:117–124.
- McGowan SJ, Holden MT, Bycroft BW, Salmond GP. Molecular genetics of carbapenem antibiotic biosynthesis. *Antonie Van Leeuwenhoek* 1999, 75:135–141.
- Minogue TD, Carlier AL, Koutsoudis MD, von Bodman SB. The cell density-dependent expression of stewartia exopolysaccharide in *Pantoea stewartii* ssp. *Stewartii* is a function of EsaR-mediated repression of the *rcaA* gene. *Mol Microbiol* 2005, 56:189–203.
- Hilton T, Rosche T, Froelich B, Smith B, Oliver J. Capsular polysaccharide phase variation in *Vibrio vulnificus*. *Appl Environ Microbiol* 2006, 72:6986–6993.
- Koutsoudis MD, Tsaltas D, Minogue TD, von Bodman SB. Quorum-sensing regulation governs bacterial adhesion, biofilm development, and host colonization in *Pantoea stewartii* subspecies *stewartii*. *Proc Natl Acad Sci U S A* 2006, 103:5983–5988.
- Dunlap PV. Quorum regulation of luminescence in *Vibrio fischeri*. *J Mol Microbiol Biotechnol* 1999, 1:5–12.
- Nyholm SV, McFall-Ngai MJ. The winnowing: establishing the squid-vibrio symbiosis. *Nat Rev Microbiol* 2004, 2:632–642.
- Daniels R, Vanderleyden J, Michiels J. Quorum sensing and swarming migration in bacteria. *FEMS Microbiol Rev* 2004, 28:261–289.
- Atkinson S, Chang CY, Sockett RE, Camara M, Williams P. Quorum sensing in *Yersinia enterocolitica* controls swimming and swarming motility. *J Appl Bacteriol* 2006, 188:1451–1461.
- Hussain MB, Zhang HB, Xu JL, Liu Q, Jiang Z, et al. The acyl-homoserine lactone-type quorum-sensing system modulates cell motility and virulence of *Erwinia chrysanthemi* pv. *zeae*. *J Appl Bacteriol* 2008, 190:1045–1053.
- Dunny GM, Winans SC. *Cell-Cell Signaling in Bacteria*. Washington, D.C: ASM Press; 1999.
- Nealson KH, Platt T, Hastings JW. Cellular control of the synthesis and activity of the bacterial luminescent system. *J Appl Bacteriol* 1970, 104:313–322.
- Engbrecht J, Silverman M. Identification of genes and gene products necessary for bacterial bioluminescence. *Proc Natl Acad Sci U S A* 1984, 81:4154–4158.
- Fuqua WC, Winans SC, Greenberg EP. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J Appl Bacteriol* 1994, 176:269–275.

16. Fuqua C, Winans SC, Greenberg EP. Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu Rev Microbiol* 1996, 50:727–751.
17. Swift S, Throup JP, Williams P, Salmond GP, Stewart GS. Quorum sensing: a population-density component in the determination of bacterial phenotype. *Trends Biochem Sci* 1996, 21:214–219.
18. Swift S, Stewart GS, Williams P. The inner workings of a quorum sensing signal generator. *Trends Microbiol* 1996, 4:463–465.
19. Surette MG, Bassler BL. Quorum sensing in *Escherichia coli* and *Salmonella typhimurium*. *Proc Natl Acad Sci U S A* 1998, 95:7046–7050.
20. Hardman AM, Stewart GS, Williams P. Quorum sensing and the cell-cell communication dependent regulation of gene expression in pathogenic and non-pathogenic bacteria. *Antonie Van Leeuwenhoek* 1998, 74:199–210.
21. Surette MG, Miller MB, Bassler BL. Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: a new family of genes responsible for autoinducer production. *Proc Natl Acad Sci U S A* 1999, 96:1639–1644.
22. Parsek MR, Val DL, Hanzelka BL, Cronan JE Jr, Greenberg EP. Acyl homoserine-lactone quorum-sensing signal generation. *Proc Natl Acad Sci USA* 1999, 96:4360–4365.
23. Eberl L. N-acyl homoserinylactone-mediated gene regulation in gram-negative bacteria. *Syst Appl Microbiol* 1999, 22:493–506.
24. Williams P, Camara M, Hardman A, Swift S, Milton D, et al. Quorum sensing and the population-dependent control of virulence. *Philos Trans R Soc Lond B Biol Sci* 2000, 355:667–680.
25. Parsek MR, Greenberg EP. Acyl-homoserine lactone quorum sensing in gram-negative bacteria: a signaling mechanism involved in associations with higher organisms. *Proc Natl Acad Sci U S A* 2000, 97:8789–8793.
26. Miller MB, Bassler BL. Quorum sensing in bacteria. *Annu Rev Microbiol* 2001, 55:165–199.
27. Fuqua C, Parsek MR, Greenberg EP. Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu Rev Genet* 2001, 35:439–468.
28. Bassler BL. Small talk. Cell-to-cell communication in bacteria. *Cell* 2002, 109:421–424.
29. von Bodman SB, Bauer WD, Coplin DL. Quorum sensing in plant-pathogenic bacteria. *Annu Rev Phytopathol* 2003, 41:455–482.
30. Xavier KB, Bassler BL. LuxS quorum sensing: more than just a numbers game. *Curr Opin Microbiol* 2003, 6:191–197.
31. Taga ME, Bassler BL. Chemical communication among bacteria. *Proc Natl Acad Sci U S A* 2003, 100(suppl 2):14549–14554.
32. Henke JM, Bassler BL. Bacterial social engagements. *Trends Cell Biol* 2004, 14:648–656.
33. Waters CM, Bassler BL. Quorum sensing: cell-to-cell communication in bacteria. *Annu Rev Cell Dev Biol* 2005, 21:319–346.
34. Camilli A, Bassler BL. Bacterial small-molecule signaling pathways. *Science* 2006, 311:1113–1116.
35. Milton DL. Quorum sensing in vibrios: complexity for diversification. *Int J Med Microbiol* 2006, 296:61–71.
36. Williams P, Winzer K, Chan WC, Camara M. Look who's talking: communication and quorum sensing in the bacterial world. *Philos Trans R Soc Lond B Biol Sci* 2007, 362:1119–1134.
37. Dunny GM. The peptide pheromone-inducible conjugation system of *Enterococcus faecalis* plasmid pCF10: cell-cell signalling, gene transfer, complexity and evolution. *Philos Trans R Soc Lond B Biol Sci* 2007, 362:1185–1193.
38. Williams P. Quorum sensing, communication and cross-kingdom signalling in the bacterial world. *Microbiology* 2007, 153:3923–3938.
39. Dunny GM, Brickman TJ, Dworkin M. Multicellular behavior in bacteria: communication, cooperation, competition and cheating. *Bioessays* 2008, 30:296–298.
40. Hughes DT, Sperandio V. Inter-kingdom signalling: communication between bacteria and their hosts. *Nat Rev Microbiol* 2008, 6:111–120.
41. West SA, Griffin AS, Gardner A, Diggle SP. Social evolution theory for microorganisms. *Nat Rev Microbiol* 2006, 4:597–607.
42. Diggle SP, Griffin AS, Campbell GS, West SA. Cooperation and conflict in quorum-sensing bacterial populations. *Nature* 2007, 450:411–414.
43. Finch RG, Pritchard DI, Bycroft BW, Williams P, Stewart GS. Quorum sensing: a novel target for anti-infective therapy. *J Antimicrob Chemother* 1998, 42:569–571.
44. Dong YH, Wang LH, Xu JL, Zhang HB, Zhang XF, et al. Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature* 2001, 411:813–817.
45. Zhang LH. Quorum quenching and proactive host defense. *Trends Plant Sci* 2003, 8:238–244.
46. Rice SA, McDougald D, Kumar N, Kjelleberg S. The use of quorum-sensing blockers as therapeutic agents for the control of biofilm-associated infections. *Curr Opin Investig Drugs* 2005, 6:178–184.
47. Rasmussen TB, Givskov M. Quorum sensing inhibitors: a bargain of effects. *Microbiology* 2006, 152:895–904.

48. Rasmussen TB, Givskov M. Quorum-sensing inhibitors as anti-pathogenic drugs. *Int J Med Microbiol* 2006, 296:149–161.
49. Chan WC, Coyle BJ, Williams P. Virulence regulation and quorum sensing in staphylococcal infections: competitive AgrC antagonists as quorum sensing inhibitors. *J Med Chem* 2004, 47:4633–4641.
50. Dong YH, Wang LY, Zhang LH. Quorum-quenching microbial infections: mechanisms and implications. *Philos Trans R Soc Lond B Biol Sci* 2007, 362:1201–1211.
51. Gottesman S. Bacterial regulation: global regulatory networks. *Annu Rev Genet* 1984, 18:415–441.
52. VanBogelen RA, Greis KD, Blumenthal RM, Tani TH, Matthews RG. Mapping regulatory networks in microbial cells. *Trends Microbiol* 1999, 7:320–328.
53. Gilman A, Arkin AP. Genetic “code”: representations and dynamical models of genetic components and networks. *Annu Rev Genomics Hum Genet* 2002, 3:3341–3369.
54. Herrgard MJ, Covert MW, Palsson BO. Reconstruction of microbial transcriptional regulatory networks. *Curr Opin Biotechnol* 2004, 15:70–77.
55. Fuqua C, Greenberg EP. Self perception in bacteria: quorum sensing with acylated homoserine lactones. *Curr Opin Microbiol* 1998, 1:183–189.
56. Whitehead NA, Barnard AM, Slater H, Simpson NJ, Salmond GP. Quorum-sensing in Gram-negative bacteria. *FEMS Microbiol Rev* 2001, 25:365–404.
57. Kleerebezem M, Quadri LE, Kuipers OP, de Vos WM. Quorum sensing by peptide pheromones and two-component signal-transduction systems in Gram-positive bacteria. *Mol Microbiol* 1997, 24:895–904.
58. Dunny GM, Leonard BA. Cell-cell communication in gram-positive bacteria. *Annu Rev Microbiol* 1997, 51:527–564.
59. Lazazzera BA, Grossman AD. The ins and outs of peptide signaling. *Trends Microbiol* 1998, 6:288–294.
60. Lyon GJ, Novick RP. Peptide signaling in *Staphylococcus aureus* and other Gram-positive bacteria. *Peptides* 2004, 25:1389–1403.
61. Podbielski A, Kreikemeyer B. Cell density—dependent regulation: basic principles and effects on the virulence of Gram-positive cocci. *Int J Infect Dis* 2004, 8:81–95.
62. Pottathil M, Lazazzera BA. The extracellular Phr peptide-Rap phosphatase signaling circuit of *Bacillus subtilis*. *Front Biosci* 2003, 8:d32–d45.
63. Leonard BA, Podbielski A, Hedberg PJ, Dunny GM. Enterococcus faecalis pheromone binding protein, PrgZ, recruits a chromosomal oligopeptide permease system to import sex pheromone cCF10 for induction of conjugation. *Proc Natl Acad Sci U S A* 1996, 93:260–264.
64. Pearson JP, van Delden C, Iglewski BH. Active efflux and diffusion are involved in transport of *Pseudomonas aeruginosa* cell-to-cell signals. *J Appl Bacteriol* 1999, 181:1203–1210.
65. Gray KM, Garey JR. The evolution of bacterial LuxI and LuxR quorum sensing regulators. *Microbiology* 2001, 147:2379–2387.
66. Lerat E, Moran NA. The evolutionary history of quorum-sensing systems in bacteria. *Mol Biol Evol* 2004, 21:903–913.
67. Gilson L, Kuo A, Dunlap PV. AinS and a new family of autoinducer synthesis proteins. *J Appl Bacteriol* 1995, 177:6946–6951.
68. Ortori CA, Atkinson S, Chhabra SR, Camara M, Williams P, et al. Comprehensive profiling of N-acylhomoserine lactones produced by *Yersinia pseudotuberculosis* using liquid chromatography coupled to hybrid quadrupole-linear ion trap mass spectrometry. *Anal Bioanal Chem* 2007, 387:497–511.
69. Kobayashi H, Kaern M, Araki M, Chung K, Gardner TS, et al. Programmable cells: interfacing natural and engineered gene networks. *Proc Natl Acad Sci U S A* 2004, 101:8414–8419.
70. Wang C, Zhang HB, Wang LH, Zhang LH. Succinic semialdehyde couples stress response to quorum-sensing signal decay in *Agrobacterium tumefaciens*. *Mol Microbiol* 2006, 62:45–56.
71. Lazazzera BA. The intracellular function of extracellular signaling peptides. *Peptides* 2001, 22:1519–1527.
72. Perego M, Brannigan JA. Pentapeptide regulation of aspartyl-phosphate phosphatases. *Peptides* 2001, 22:1541–1547.
73. Schauder S, Shokat K, Surette MG, Bassler BL. The LuxS family of bacterial autoinducers: biosynthesis of a novel quorum-sensing signal molecule. *Mol Microbiol* 2001, 41:463–476.
74. Vendeville A, Winzer K, Heurlier K, Tang CM, Hardie KR. Making ‘sense’ of metabolism: autoinducer-2, LuxS and pathogenic bacteria. *Nat Rev Microbiol* 2005, 3:383–396.
75. Neiditch MB, Federle MJ, Miller ST, Bassler BL, Hughson FM. Regulation of LuxPQ receptor activity by the quorum-sensing signal autoinducer-2. *Mol Cell* 2005, 18:507–518.
76. Xavier KB, Miller ST, Lu W, Kim JH, Rabinowitz J, et al. Phosphorylation and processing of the quorum-sensing molecule autoinducer-2 in enteric bacteria. *ACS Chem Biol* 2007, 2:128–136.
77. Diggle SP, Cornelis P, Williams P, Camara M. 4-quinolone signalling in *Pseudomonas aeruginosa*: old molecules, new perspectives. *Int J Med Microbiol* 2006, 296:83–91.

78. Pesci EC, Milbank JB, Pearson JP, McKnight S, Kende AS, et al. Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 1999, 96:11229–11234.
79. Fouhy Y, Lucey JF, Ryan RP, Dow JM. Cell-cell signaling, cyclic di-GMP turnover and regulation of virulence in *Xanthomonas campestris*. *Res Microbiol* 2006, 157:899–904.
80. Van Houdt R, Givskov M, Michiels CW. Quorum sensing in *Serratia*. *FEMS Microbiol Rev* 2007, 31:407–424.
81. Higgins DA, Pomianek ME, Kraml CM, Taylor RK, Semmelhack MF, et al. The major *Vibrio cholerae* autoinducer and its role in virulence factor production. *Nature* 2007, 450:883–886.
82. Takano E. Gamma-butyrolactones: Streptomyces signalling molecules regulating antibiotic production and differentiation. *Curr Opin Microbiol* 2006, 9:287–294.
83. Nasser W, Reverchon S. New insights into the regulatory mechanisms of the LuxR family of quorum sensing regulators. *Anal Bioanal Chem* 2007, 387:381–390.
84. Urbanowski ML, Lostroh CP, Greenberg EP. Reversible acyl-homoserine lactone binding to purified *Vibrio fischeri* LuxR protein. *J Appl Bacteriol* 2004, 186:631–637.
85. Zhu J, Winans SC. Autoinducer binding by the quorum-sensing regulator TraR increases affinity for target promoters in vitro and decreases TraR turnover rates in whole cells. *Proc Natl Acad Sci U S A* 1999, 96:4832–4837.
86. Zhu J, Winans SC. The quorum-sensing transcriptional regulator TraR requires its cognate signaling ligand for protein folding, protease resistance, and dimerization. *Proc Natl Acad Sci U S A* 2001, 98:1507–1512.
87. McGowan SJ, Barnard AM, Bosgelmez G, Sebahia M, Simpson NJ, et al. Carbapenem antibiotic biosynthesis in *Erwinia carotovora* is regulated by physiological and genetic factors modulating the quorum sensing-dependent control pathway. *Mol Microbiol* 2005, 55:526–545.
88. von Bodman SB, Majerczak DR, Coplin DL. A negative regulator mediates quorum-sensing control of exopolysaccharide production in *Pantoea stewartii* subsp. *stewartii*. *Proc Natl Acad Sci U S A* 1998, 95:7687–7692.
89. Andersson RA, Eriksson AR, Heikinheimo R, Mae A, Pirhonen M, et al. Quorum sensing in the plant pathogen *Erwinia carotovora* subsp. *carotovora*: the role of expR(Ecc). *Mol Plant Microbe Interact* 2000, 13:384–393.
90. Minogue TD, Wehland-von Trebra M, Bernhard F, von Bodman SB. The autoregulatory role of EsaR, a quorum-sensing regulator in *Pantoea stewartii* ssp. *stewartii*: evidence for a repressor function. *Mol Microbiol* 2002, 44:1625–1635.
91. Burr T, Barnard AM, Corbett MJ, Pemberton CL, Simpson NJ, et al. Identification of the central quorum sensing regulator of virulence in the enteric phytopathogen, *Erwinia carotovora*: the VirR repressor. *Mol Microbiol* 2006, 59:113–125.
92. Nasser W, Bouillant ML, Salmond G, Reverchon S. Characterization of the *Erwinia chrysanthemi* expI-expR locus directing the synthesis of two N-acyl-homoserine lactone signal molecules. *Mol Microbiol* 1998, 29:1391–1405.
93. Sjoblom S, Brader G, Koch G, Palva ET. Cooperation of two distinct ExpR regulators controls quorum sensing specificity and virulence in the plant pathogen *Erwinia carotovora*. *Mol Microbiol* 2006, 60:1474–1489.
94. Cui Y, Chatterjee A, Hasegawa H, Chatterjee AK. *Erwinia carotovora* subspecies produce duplicate variants of ExpR, LuxR homologs that activate rsmA transcription but differ in their interactions with N-acylhomoserine lactone signals. *J Appl Bacteriol* 2006, 188:4715–4726.
95. Shadel GS, Baldwin TO. Positive autoregulation of the *Vibrio fischeri* luxR gene. LuxR and autoinducer activate cAMP-catabolite gene activator protein complex-independent and -dependent luxR transcription. *J Biochem Mol Biol Biophys* 1992, 267:7696–7702.
96. Atkinson S, Sockett RE, Camara M, Williams P. Quorum sensing and the lifestyle of *Yersinia*. *Curr Issues Mol Biol* 2006, 8:1–10.
97. Atkinson S, Chang CY, Patrick HL, Buckley CM, Wang Y, et al. Functional interplay between the *Yersinia pseudotuberculosis* YpsRI and YtbRI quorum sensing systems modulates swimming motility by controlling expression of flhDC and fliA. *Mol Microbiol* 2008, 69:137–151.
98. White CE, Winans SC. Cell-cell communication in the plant pathogen *Agrobacterium tumefaciens*. *Philos Trans R Soc Lond B Biol Sci* 2007, 362:1135–1148.
99. Shadel GS, Baldwin TO. The *Vibrio fischeri* LuxR protein is capable of bidirectional stimulation of transcription and both positive and negative regulation of the luxR gene. *J Appl Bacteriol* 1991, 173:568–574.
100. Schuster M, Greenberg EP. A network of networks: quorum-sensing gene regulation in *Pseudomonas aeruginosa*. *Int J Med Microbiol* 2006, 296:73–81.
101. Showalter RE, Martin MO, Silverman MR. Cloning and nucleotide sequence of luxR, a regulatory gene controlling bioluminescence in *Vibrio harveyi*. *J Appl Bacteriol* 1990, 172:2946–2954.
102. Jobling MG, Holmes RK. Characterization of hapR, a positive regulator of the *Vibrio cholerae* HA/protease gene hap, and its identification as a functional homologue of the *Vibrio harveyi* luxR gene. *Mol Microbiol* 1997, 26:1023–1034.

103. Hammer BK, Bassler BL. Quorum sensing controls biofilm formation in *Vibrio cholerae*. *Mol Microbiol* 2003, 50:101–104.
104. Zhu J, Miller MB, Vance RE, Dziejman M, Bassler BL, et al. Quorum-sensing regulators control virulence gene expression in *Vibrio cholerae*. *Proc Natl Acad Sci U S A* 2002, 99:3129–3134.
105. Fidopiastis PM, Miyamoto CM, Jobling MG, Meighen EA, Ruby EG. LitR, a new transcriptional activator in *Vibrio fischeri*, regulates luminescence and symbiotic light organ colonization. *Mol Microbiol* 2002, 45:131–143.
106. De Silva RS, Kovacicova G, Lin W, Taylor RK, Skorpinski K, et al. Crystal structure of the *Vibrio cholerae* quorum-sensing regulatory protein HapR. *J Appl Bacteriol* 2007, 189:5683–5691.
107. Ramos JL, Martinez-Bueno M, Molina-Henares AJ, Teran W, Watanabe K, et al. The TetR family of transcriptional repressors. *Microbiol Mol Biol Rev* 2005, 69:326–356.
108. Meibom KL, Blokesch M, Dolganov NA, Wu CY, Schoolnik GK. Chitin induces natural competence in *Vibrio cholerae*. *Science* 2005, 310:1824–1827.
109. Lin YH, Miyamoto C, Meighen EA. Purification and characterization of a luxO promoter binding protein LuxT from *Vibrio harveyi*. *Protein Expr Purif* 2000, 20:87–94.
110. Lenz DH, Mok KC, Lilley BN, Kulkarni RV, Wingreen NS, et al. The small RNA chaperone Hfq and multiple small RNAs control quorum sensing in *Vibrio harveyi* and *Vibrio cholerae*. *Cell* 2004, 118:69–82.
111. Henke JM, Bassler BL. Three parallel quorum-sensing systems regulate gene expression in *Vibrio harveyi*. *J Appl Bacteriol* 2004, 186:6902–6914.
112. Lenz DH, Miller MB, Zhu J, Kulkarni RV, Bassler BL. CsrA and three redundant small RNAs regulate quorum sensing in *Vibrio cholerae*. *Mol Microbiol* 2005, 58:1186–1202.
113. Yarwood JM, Schlievert PM. Quorum sensing in Staphylococcus infections. *J Clin Invest* 2003, 112:1620–1625.
114. Tyson JJ, Chen KC, Novak B. Sniffers, buzzers, toggles and blinkers: dynamics of regulatory and signaling pathways in the cell. *Curr Opin Cell Biol* 2003, 15:221–231.
115. Ferrell JE Jr. Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability. *Curr Opin Cell Biol* 2002, 14:140–148.
116. Ozbudak EM, Thattai M, Lim HN, Shraiman BI, Van Oudenaarden A. Multistability in the lactose utilization network of *Escherichia coli*. *Nature* 2004, 427:737–740.
117. Santillan M, Mackey MC. Quantitative approaches to the study of bistability in the lac operon of *Escherichia coli*. *J R Soc Interface* 2008, 5(suppl 1):S29–S39.
118. Tyson JJ, Chen K, Novak B. Network dynamics and cell physiology. *Nat Rev Mol Cell Biol* 2001, 2:908–916.
119. Dockery JD, Keener JP. A mathematical model for quorum sensing in *Pseudomonas aeruginosa*. *Bull Math Biol* 2001, 63:95–116.
120. James S, Nilsson P, James G, Kjelleberg S, Fagerstrom T. Luminescence control in the marine bacterium *Vibrio fischeri*: an analysis of the dynamics of lux regulation. *J Biochem Mol Biol* 2000, 296:1127–1137.
121. Goryachev AB, Toh DJ, Lee T. Systems analysis of a quorum sensing network: design constraints imposed by the functional requirements, network topology and kinetic constants. *Biosystems* 2006, 83:178–187.
122. Kaufmann BB, Van Oudenaarden A. Stochastic gene expression: from single molecules to the proteome. *Curr Opin Genet Dev* 2007, 17:107–112.
123. Elowitz MB, Levine AJ, Siggia ED, Swain PS. Stochastic gene expression in a single cell. *Science* 2002, 297:1183–1186.
124. Welch M, Todd DE, Whitehead NA, McGowan SJ, Bycroft BW, et al. N-acyl homoserine lactone binding to the CarR receptor determines quorum-sensing specificity in *Erwinia*. *EMBO J* 2000, 19:631–641.
125. Zhu J, Oger PM, Schrammeijer B, Hooykaas PJ, Farand SK, et al. The bases of crown gall tumorigenesis. *J Appl Bacteriol* 2000, 182:3885–3895.
126. Zhang RG, Pappas T, Brace JL, Miller PC, Oulmassov T, et al. Structure of a bacterial quorum-sensing transcription factor complexed with pheromone and DNA. *Nature* 2002, 417:971–974.
127. Brennan RG, Link TM. Hfq structure, function and ligand binding. *Curr Opin Microbiol* 2007, 10:125–133.
128. Salazar C, Hofer T. Competition effects shape the response sensitivity and kinetics of phosphorylation cycles in cell signaling. *Ann N Y Acad Sci* 2006, 109:1517–1530.
129. Neves SR, Iyengar R. Modeling of signaling networks. *Bioessays* 2002, 24:1110–1117.
130. Vandenesch F, Kornblum J, Novick RP. A temporal signal, independent of agr, is required for hla but not spa transcription in *Staphylococcus aureus*. *J Appl Bacteriol* 1991, 173:6313–6320.
131. Majdalani N, Vanderpool CK, Gottesman S. Bacterial small RNA regulators. *Crit Rev Biochem Mol Biol* 2005, 40:93–113.
132. Babitzke P, Romeo T. CsrB sRNA family: sequestration of RNA-binding regulatory proteins. *Curr Opin Microbiol* 2007, 10:156–163.

133. Kay E, Dubuis C, Haas D. Three small RNAs jointly ensure secondary metabolism and biocontrol in *Pseudomonas fluorescens* CHA0. *Proc Natl Acad Sci U S A* 2005, 102:17136–17141.
134. Levine E, Zhang Z, Kuhlman T, Hwa T. Quantitative characteristics of gene regulation by small RNA. *PLoS Biol* 2007, 5:e229.
135. Aiba H. Mechanism of RNA silencing by Hfq-binding small RNAs. *Curr Opin Microbiol* 2007, 10:134–139.
136. Shimoni Y, Friedlander G, Hetzroni G, Niv G, Altuvia S, et al. Regulation of gene expression by small non-coding RNAs: a quantitative view. *Mol Syst Biol* 2007, 3:138.
137. Bejerano-Sagie M, Xavier KB. The role of small RNAs in quorum sensing. *Curr Opin Microbiol* 2007, 10:189–198.
138. Barnard AM, Salmond GP. Quorum sensing in *Erwinia* species. *Anal Bioanal Chem* 2007, 387:415–423.
139. Hwang I, Smyth AJ, Luo ZQ, Farrand SK. Modulating quorum sensing by antiactivation: TraM interacts with TraR to inhibit activation of Ti plasmid conjugal transfer genes. *Mol Microbiol* 1999, 34:282–294.
140. Swiderska A, Berndtson AK, Cha MR, Li L, Beaudoin GM III, et al. Inhibition of the *Agrobacterium tumefaciens* TraR quorum-sensing regulator. Interactions with the TraM anti-activator. *J Biochem Mol Biol Biophys* 2001, 276:49449–49458.
141. Qin Y, Su S, Farrand SK. Molecular basis of transcriptional antiactivation. TraM disrupts the TraR-DNA complex through stepwise interactions. *J Biochem Mol Biol Biophys* 2007, 282:19979–19991.
142. Goryachev AB, Toh DJ, Wee KB, Lee T, Zhang HB, et al. Transition to quorum sensing in an *Agrobacterium* population: a stochastic model. *PLoS Comput Biol* 2005, 1:e37.
143. Withers H, Swift S, Williams P. Quorum sensing as an integral component of gene regulatory networks in Gram-negative bacteria. *Curr Opin Microbiol* 2001, 4:186–193.
144. Venturi V. Regulation of quorum sensing in *Pseudomonas*. *FEMS Microbiol Rev* 2006, 30:274–291.
145. Juhas M, Eberl L, Tummeler B. Quorum sensing: the power of cooperation in the world of *Pseudomonas*. *Environ Microbiol* 2005, 7:459–471.
146. Milton DL, Chalker VJ, Kirke D, Hardman A, Camara M, et al. The LuxM homologue VanM from *Vibrio anguillarum* directs the synthesis of N-(3-hydroxyhexanoyl)homoserine lactone and N-hexanoylhomoserine lactone. *J Appl Bacteriol* 2001, 183:3537–3547.
147. Miller MB, Skorupski K, Lenz DH, Taylor RK, Bassler BL. Parallel quorum sensing systems converge to regulate virulence in *Vibrio cholerae*. *Cell* 2002, 110:303–314.
148. Waters CM, Bassler BL. The *Vibrio harveyi* quorum-sensing system uses shared regulatory components to discriminate between multiple autoinducers. *Genes Dev* 2006, 20:2754–2767.
149. Latifi A, Foglino M, Tanaka K, Williams P, Lazdunski A. A hierarchical quorum-sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators LasR and RhIR (VsmR) to expression of the stationary-phase sigma factor RpoS. *Mol Microbiol* 1996, 21:1137–1146.
150. Pesci EC, Pearson JP, Seed PC, Iglewski BH. Regulation of las and rhl quorum sensing in *Pseudomonas aeruginosa*. *J Appl Bacteriol* 1997, 179:3127–3132.
151. Lupp C, Urbanowski M, Greenberg EP, Ruby EG. The *Vibrio fischeri* quorum-sensing systems ain and lux sequentially induce luminescence gene expression and are important for persistence in the squid host. *Mol Microbiol* 2003, 50:319–331.
152. Lupp C, Ruby EG. *Vibrio fischeri* uses two quorum-sensing systems for the regulation of early and late colonization factors. *J Appl Bacteriol* 2005, 187:3620–3629.
153. Antunes LC, Schaefer AL, Ferreira RB, Qin N, Stevens AM, et al. Transcriptome analysis of the *Vibrio fischeri* LuxR-LuxI regulon. *J Appl Bacteriol* 2007, 189:8387–8391.
154. Majdalani N, Gottesman S. The Rcs phosphorelay: a complex signal transduction system. *Annu Rev Microbiol* 2005, 59:379–405.
155. Wehland M, Kiecker C, Coplin DL, Kelm O, Saenger W, et al. Identification of an RcsA/RcsB recognition motif in the promoters of exopolysaccharide biosynthetic operons from *Erwinia amylovora* and *Pantoea stewartii* subsp. *stewartii*. *J Biochem Mol Biol Biophys* 1999, 274:3300–3307.
156. Carlier AL, von Bodman SB. The rcsA promoter of *Pantoea stewartii* subsp. *stewartii* features a low-level constitutive promoter and an EsaR quorum-sensing-regulated promoter. *J Appl Bacteriol* 2006, 188:4581–4584.
157. Romling U, Amikam D. Cyclic di-GMP as a second messenger. *Curr Opin Microbiol* 2006, 9:218–228.
158. Jenal U, Malone J. Mechanisms of cyclic-di-GMP signaling in bacteria. *Annu Rev Genet* 2006, 40:385–407.
159. Ryan RP, Fouhy Y, Lucey JF, Dow JM. Cyclic di-GMP signaling in bacteria: recent advances and new puzzles. *J Appl Bacteriol* 2006, 188:8327–8334.
160. Cotter PA, Stibitz S. c-di-GMP-mediated regulation of virulence and biofilm formation. *Curr Opin Microbiol* 2007, 10:17–23.
161. Waters CM, Lu W, Rabinowitz JD, Bassler BL. Quorum sensing controls biofilm formation in *Vibrio*

- cholerae* through modulation of cyclic di-GMP levels and repression of vpsT. *J Appl Bacteriol* 2008, 190:2527–2536.
162. Rahman M, Simm R, Kader A, Basseres E, Romling U, et al. The role of c-di-GMP signaling in an *Aeromonas veronii* biovar sobria strain. *FEMS Microbiol Lett* 2007, 273:172–179.
  163. Heermann R, Fuchs TM. Comparative analysis of the *Photobacterium luminescens* and the *Yersinia enterocolitica* genomes: uncovering candidate genes involved in insect pathogenicity. *BMC Genomics* 2008, 9:40.
  164. Williamson NR, Fineran PC, Ogawa W, Woodley LR, Salmond GP. Integrated regulation involving quorum sensing, a two-component system, a GGDEF/EAL domain protein and a post-transcriptional regulator controls swarming and RhlA-dependent surfactant biosynthesis in *Serratia*. *Environ Microbiol* 2008, 10:1202–1217.
  165. Koerber AJ, King JR, Williams P. Deterministic and stochastic modelling of endosome escape by *Staphylococcus aureus*: “quorum” sensing by a single bacterium. *J Math Biol* 2005, 50:440–488.
  166. Redfield RJ. Is quorum sensing a side effect of diffusion sensing? *Trends Microbiol* 2002, 10:365–370.
  167. McLean RJ, Whiteley M, Stickler DJ, Fuqua WC. Evidence of autoinducer activity in naturally occurring biofilms. *FEMS Microbiol Lett* 1997, 154:259–263.
  168. Stickler DJ, Morris NS, McLean RJ, Fuqua C. Biofilms on indwelling urethral catheters produce quorum-sensing signal molecules in situ and in vitro. *Appl Environ Microbiol* 1998, 64:3486–3490.
  169. Parsek MR, Greenberg EP. Quorum sensing signals in development of *Pseudomonas aeruginosa* biofilms. *Methods Enzymol* 1999, 310:43–55.
  170. Singh PK, Schaefer AL, Parsek MR, Moninger TO, Welsh MJ, et al. Quorum-sensing signals indicate that cystic fibrosis lungs are infected with bacterial biofilms. *Nature* 2000, 407:762–764.
  171. Schembri MA, Givskov M, Klemm P. An attractive surface: gram-negative bacterial biofilms. *Sci STKE* 2002, 132:RE6.
  172. Chopp DL, Kirisits MJ, Moran B, Parsek MR. A mathematical model of quorum sensing in a growing bacterial biofilm. *J Ind Microbiol Biotechnol* 2002, 29:339–346.
  173. Ward JP, King JR, Koerber AJ, Croft JM, Sockett RE, et al. Early development and quorum sensing in bacterial biofilms. *J Math Biol* 2003, 47:23–55.
  174. Lenz AP, Williamson K, Pitts B, Stewart PS, Franklin MJ. Localized gene expression in *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol* 2008, 74:4463–4471.
  175. Wagner VE, Iglewski BH. *P. aeruginosa* biofilms in CF infection. *Clin Rev Allergy Immunol* 2008, 35:124–134.
  176. Boles BR, Horswill AR. Agr-mediated dispersal of *Staphylococcus aureus* biofilms. *PLoS Pathog* 2008, 4:e1000052.
  177. Nadell CD, Xavier JB, Levin SA, Foster KR. The evolution of quorum sensing in bacterial biofilms. *PLoS Biol* 2008, 6:e14.
  178. Losick R, Kaiser D. Why and how bacteria communicate. *Sci Am* 1997, 276:68–73.
  179. Dworkin M. Shape, polarity, and multicellular morphogenesis. *Curr Opin Microbiol* 2007, 10:588–590.
  180. Aguilar C, Vlamakis H, Losick R, Kolter R. Thinking about *Bacillus subtilis* as a multicellular organism. *Curr Opin Microbiol* 2007, 10:638–643.
  181. Kaiser D. Coupling cell movement to multicellular development in myxobacteria. *Nat Rev Microbiol* 2003, 1:45–54.
  182. Kholodenko BN. Cell-signalling dynamics in time and space. *Nat Rev Mol Cell Biol* 2006, 7:165–176.
  183. Viretta AU, Fussenegger M. Modeling the quorum sensing regulatory network of human-pathogenic *Pseudomonas aeruginosa*. *Biotechnol Prog* 2004, 20:670–678.
  184. Ward JP, King JR, Koerber AJ, Williams P, Croft JM, et al. Mathematical modelling of quorum sensing in bacteria. *IMA J Math Appl Med Biol* 2001, 18:263–292.

## FURTHER READING

Quorum sensing represents an example of bacterial cell–cell communication. A generally accessible and concise introduction to this topic can be found in the enlightening and visually appealing review<sup>178</sup> by Losick and Kaiser. A systematic account of this field, including quorum sensing, is compiled in the comprehensive collection edited by Dunny and Winans.<sup>12</sup> Because communication is crucial for complex organization, bacterial multicellular morphogenesis is another exciting topic related to quorum sensing.<sup>179–181</sup> Those interested in application of mathematical modeling to the analysis of QSN are referred to the clear and accessible paper by Dockery and Keener.<sup>119</sup> Further references on modeling and its role in understanding intracellular decision circuitry can be found in the excellent reviews written by the pioneers of this field.<sup>114,115,118,182</sup>